

SOME REMARKS ON A GENERAL CONSTRUCTION OF APPROXIMATION PROCESSES

Lorenzo D'Ambrosio, Elisabetta Mangino
Ref. SISSA 8/2001/M

The aim of this paper is to study the characteristics of a general method to produce a new approximation sequence from a given one, by using suitable convex combinations, whose coefficients depend on some functions λ_n ($n \geq 1$). This type of construction was first used to generate Lototsky-Schnabl operators from Bernstein operators (see section 6.1. in [1]) and turned to be really useful in the study of evolution equations by means of positive operators. More precisely, the new sequences of operators were used to represent the solutions of degenerate elliptic parabolic equations, thus providing several qualitative informations both on the solutions of the equation and on the Markov process associated with the equation (see Chapter 6 in [1] and the references quoted therein and e.g. [2, 3, 4]).

In the present note we investigate which properties of the original sequence of operators $(L_n)_{n \geq 1}$ are inherited by the new sequence $(L_{n,\lambda_n})_{n \geq 1}$. More precisely, approximation properties and rates of approximations are studied in the first section. The second section is devoted to the study of the asymptotic behaviour of the sequence $(L_{n,\lambda_n})_{n \geq 1}$ and a Voronovskaja-type relation is established. Finally in the third section the behaviour of the operators L_{n,λ_n} , when acting on convex, monotone or Hölder continuous functions is studied.

Throughout the paper, I will denote an interval on the real line and J a subinterval of I . Let $F(I)$ be the space of all real valued functions on I and let $C(I)$ denote the space of all real valued continuous functions on I . Moreover set $C_b(I)$ for be the subspace of all bounded continuous functions on I .

The previous spaces of bounded functions, endowed with the natural order and the norm

$$\|f\|_\infty := \sup_{x \in I} |f(x)|, \quad f \in C_b(I),$$

are Banach lattices.

If $g : I \rightarrow \mathbb{R}$ is a strictly positive function, let $C(I, g)$ be the subspace of $C(I)$ of all functions f such that f/g is bounded.

Set $e_n(x) := x^n$ for every $x \in I$ and $n \in \mathbb{N}$, and denote by $\mathbf{1}$ the function with constant value 1. Moreover, for every $x \in I$, let $\psi_x(t) := (t - x)$ ($t \in I$). For every $f \in C(I)$, for every $\alpha \in [0, 1]$ and $x \in I$, set

$$f_{\alpha,x}(t) := f(\alpha t + (1 - \alpha)x) \quad (t \in I).$$

Let D be a sublattice of $C(I)$ such that $\mathbf{1}, e_1 \in D$ and for every $f \in D$, $x \in I$, $\alpha \in [0, 1]$, the function $f_{\alpha,x}$ belongs to D .

In this paper $(L_n)_{n \geq 1}$ will denote a sequence of linear positive operators from D into $F(J)$ such that $L_n(\mathbf{1}) = \mathbf{1}$ ($n \geq 1$). We set $L_0(f) := f|_J$ for every $f \in D$.

For every $n \geq 1$, we consider $\lambda_n : J \rightarrow [0, 1]$, and for every $f \in D$ and $x \in J$ we define the operator

$$L_{n,\lambda_n}(f)(x) := \sum_{p=0}^n \binom{n}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-p} L_p(f_{p/n,x})(x).$$

Clearly L_{n,λ_n} is a linear positive operator from D into $F(J)$.

If $\lambda_n(x) = 1$ (resp. $\lambda_n(x) = 0$) for some $x \in J$, then $L_{n,\lambda_n}(f)(x) = L_n(f)(x)$ (resp. $L_{n,\lambda_n}(f)(x) = f(x)$).

1 Approximation properties

In this section we investigate under which conditions the sequence $(L_{n,\lambda_n})_{n \geq 1}$ is an approximation process. For this we will apply some Korovkin-type theorems and their extension to spaces of unbounded functions (see [1, 6, 7, 10] and references therein).

The first step is to evaluate the operators L_{n,λ_n} on some “test” functions.

Lemma 1.1 *For every $n \geq 1$,*

$$(1.1) \quad L_{n,\lambda_n}(\mathbf{1}) = \mathbf{1}.$$

If $\psi_x^k \in D$, for $x \in J$ and $k \geq 1$, then

$$(1.2) \quad L_{n,\lambda_n}(\psi_x^k)(x) = \frac{1}{n} \sum_{p=0}^n \binom{n}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-p} \frac{p^k}{n^{k-1}} L_p(\psi_x^k)(x).$$

In particular, for any $x \in J$

$$(1.3) \quad L_{n,\lambda_n}(\psi_x)(x) = \lambda_n(x) \sum_{p=0}^{n-1} \binom{n-1}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-1-p} L_{p+1}(\psi_x)(x),$$

and, if $e_2 \in D$,

$$(1.4) \quad \begin{aligned} L_{n,\lambda_n}(\psi_x^2)(x) &= \\ &= \frac{\lambda_n(x)}{n} \sum_{p=0}^{n-1} \binom{n-1}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-1-p} (p+1) L_{p+1}(\psi_x^2)(x). \end{aligned}$$

Proof. The assertions follow by easy computations, once we observe that

$$\begin{aligned} \mathbf{1}_{\frac{p}{n},x} &= \mathbf{1}, \\ (\psi_x^k)_{\frac{p}{n},x} &= \frac{p^k}{n^k} \psi_x^k, \end{aligned}$$

for any $x \in I$, $0 \leq p \leq n$ and $k \geq 0$. □

Remark 1.2 *If $S : D \rightarrow F(J)$ is a positive linear operator and $e_2 \in D$, then for any $x \in J$, we have*

$$(1.5) \quad |S(\psi_x)(x)| \leq (1 + S(\mathbf{1})(x))\sqrt{S(\psi_x^2)(x)}.$$

Indeed, the relation

$$(1.6) \quad |\psi_x| \leq (\mathbf{1} + \psi_x^2 \delta^{-2})\delta,$$

holds for any $\delta > 0$. By applying S to (1.6) and then choosing $\delta := S(\psi_x^2)(x)$, we obtain the claim.

In the sequel we will consider some assumptions on the operators L_n . Let $(S_n)_{n \geq 1}$ be a sequence of positive linear operator from D into $F(J)$ and $M_2 : J \rightarrow \mathbb{R}_+$ be a positive function. We say that $(S_n)_{n \geq 1}$ satisfies the condition $(\mathbf{H}_2)$ with growth M_2 , if

$$(\mathbf{H}_2) \quad e_2 \in D \text{ and } S_n(\psi_x^2)(x) \leq \frac{M_2(x)}{n} \text{ for any } n \geq 1 \text{ and } x \in J.$$

If $(\mathbf{H}_2)$ is satisfied, then, by (1.5), for every $x \in J$ we have

$$(1.7) \quad \sup_{n \in \mathbb{N}} |\sqrt{n} S_n(\psi_x)(x)| \leq (1 + S_n(\mathbf{1})(x))\sqrt{M_2(x)}.$$

Sometimes we shall need of a stronger condition than (1.7). Namely, let $M_1 : J \rightarrow \mathbb{R}_+$ be a positive function, the sequence $(S_n)_{n \geq 1}$ satisfies the condition $(\mathbf{H}_1)$ with growth M_1 , if

$$(\mathbf{H}_1) \quad |S_n(\psi_x)(x)| \leq \frac{M_1(x)}{n} \text{ for any } n \geq 1 \text{ and } x \in J.$$

For instance, the conditions $(\mathbf{H}_i)$ are satisfied whenever a Voronovskaja-type relation holds (see next section).

Notice that, if for any $n \geq 1$ L_n preserves the affine functions, then from (1.1) and (1.3) we deduce that the same property holds for any L_{n, λ_n} . In this case $(\mathbf{H}_1)$ is verified with $M_1 = 0$.

The conditions $(\mathbf{H}_i)$ are preserved by the transformation $(L_n)_{n \geq 1} \rightarrow (L_{n, \lambda_n})_{n \geq 1}$, as specified by the following

Lemma 1.3 *1. If $(L_n)_{n \geq 1}$ satisfies $(\mathbf{H}_2)$ with growth M_2 , then for any $n \geq 1$ and $x \in J$ we have*

$$(1.8) \quad L_{n, \lambda_n}(\psi_x^2)(x) \leq \frac{\lambda_n(x) M_2(x)}{n},$$

therefore, also $(L_{n, \lambda_n})_{n \geq 1}$ satisfies $(\mathbf{H}_2)$ with growth M_2 , and moreover

$$(1.9) \quad |L_{n, \lambda_n}(\psi_x)(x)| \leq 2\sqrt{\frac{\lambda_n(x) M_2(x)}{n}}.$$

2. If $(L_n)_{n \geq 1}$ satisfies $(\mathbf{H}_1)$ with growth M_1 , then for any $n \geq 1$ and $x \in J$ we have

$$(1.10) \quad |L_{n, \lambda_n}(\psi_x)(x)| \leq M_1(x) \frac{1 - (1 - \lambda_n(x))^n}{n},$$

therefore, also $(L_{n, \lambda_n})_{n \geq 1}$ satisfies $(\mathbf{H}_1)$ with growth M_1 .

Proof. 1. The hypothesis $(\mathbf{H}_2)$ applied to L_p in (1.4) of Lemma 1.1 yields (1.8). Combining (1.5) and (1.8), we obtain (1.9).

2. By (1.3), we obtain

$$|L_{n,\lambda_n}(\psi_x)(x)| \leq \lambda_n(x) \sum_{p=0}^{n-1} \binom{n-1}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-1-p} \frac{M_1(x)}{p+1}.$$

Since

$$s \sum_{p=0}^{n-1} \binom{n-1}{p} \frac{s^p (1-s)^{n-1-p}}{p+1} = \frac{1}{n} \sum_{p=0}^{n-1} \binom{n}{p+1} s^{p+1} (1-s)^{n-1-p} = \frac{1 - (1-s)^n}{n},$$

holds for $0 \leq s \leq 1$, we obtain (1.10) by replacing s with $\lambda_n(x)$. $\square$

Proposition 1.4 Assume that $C_b(I) \subseteq D$ and $(L_n)_{n \geq 1}$ satisfies $(\mathbf{H}_2)$ with growth M_2 .

1. For every $f \in C_b(I)$, $n \geq 1$, $x \in J$ the following estimate holds:

$$|L_{n,\lambda_n}(f)(x) - f(x)| \leq 2\omega \left(f, \sqrt{\frac{\lambda_n(x)M_2(x)}{n}} \right).$$

2. For every differentiable $f \in D$ such that $f' \in C_b(I)$, $n \geq 1$, $x \in J$ the following estimate holds:

$$|L_{n,\lambda_n}(f)(x) - f(x)| \leq 2 \left(|f'(x)| + \omega(f', \sqrt{\frac{\lambda_n(x)M_2(x)}{n}}) \right) \sqrt{\frac{\lambda_n(x)M_2(x)}{n}},$$

and if $(\mathbf{H}_1)$ is satisfied with growth M_1 , then we have

$$\begin{aligned} |L_{n,\lambda_n}(f)(x) - f(x)| &\leq |f'(x)|M_1(x) \frac{1 - (1 - \lambda_n(x))^n}{n} + \\ &\quad + 2\omega(f', \sqrt{\frac{\lambda_n(x)M_2(x)}{n}}) \sqrt{\frac{\lambda_n(x)M_2(x)}{n}}. \end{aligned}$$

3. If $I = J = (0, \infty)$, $(L_n)_{n \geq 1}$ satisfies $(\mathbf{H}_1)$ with growth M_1 bounded and $(\mathbf{H}_2)$ with growth $M_2(x) = \alpha + \beta x + \gamma x^2$ ($x \in I$), $\alpha, \beta, \gamma \in \mathbb{R}$, and moreover we assume $L_{n,\lambda_n}(C_b(I)) \subset C_b(I)$, then for every $f \in C_b(I)$, $n \geq 1$, we have

$$\|L_{n,\lambda_n}(f) - f\|_\infty \leq K \left(\omega_{\sqrt{M_2}}^2(f, \frac{1}{\sqrt{n}}) + \frac{1}{n} \right),$$

where $K > 0$ is a constant independent of n , and

$$\omega_\phi(f, \delta) := \sup_{0 \leq h \leq \delta, x \pm h\phi(x) \geq 0} |\Delta_{h\phi(x)}^2 f(x)|$$

with

$$\Delta_{h\phi(x)}^2 f(x) = f(x - h\phi(x)) - 2f(x) + f(x + h\phi(x)).$$

If $L_n(e_1) = e_1$, the previous estimate becomes

$$\|L_{n,\lambda_n}(f) - f\|_\infty \leq \omega_{\sqrt{M_2}}^2(f, \sqrt{\frac{\|\lambda_n\|}{n}}).$$

Proof. The statements in 1. and 2. follow from Theorem 5.1.2 and the subsequent remark in [1], taking into account the estimates (1.8), (1.9) and (1.10).

3. is an immediate consequence of Theorem 1 and the subsequent remark in [12] and the estimates (1.8) and (1.10). $\square$

Remark 1.5 Note that in the setting of 3. of previous proposition, if $\|\lambda_n\| \rightarrow 0$, there also holds

$$\|L_{n,\lambda_n}(f) - f\|_\infty \leq K \left(\omega_{\sqrt{M_2}}^2(f, \sqrt{\|\lambda_n\|}) + \|\lambda_n\| \right).$$

Indeed using Bernoulli inequality $(1+s)^n \geq 1+ns$ ($s \geq -1$), from (1.10) one has $|L_{n,\lambda_n}(\psi_x)(x)| \leq M_1(x) \frac{1-(1-\lambda_n(x))^n}{n} \leq M_1(x)\lambda_n(x)$, and then, we can argue as before.

In the sequel, with $g : I \rightarrow \mathbb{R}$ we shall denote a strictly convex function such that $g \geq c$ for some constant $c > 0$.

Remark 1.6 1. If f is (strictly) convex, then also $f_{\alpha,x}$ is (strictly) convex, for any $\alpha \in [0, 1]$ and $x \in I$.

2. If $f \in C(I, g)$, then also $f_{\alpha,x} \in C(I, g)$, for any $\alpha \in [0, 1]$ and $x \in I$.

3. If $f \in C^k(I)$, then also $f_{\alpha,x} \in C^k(I)$, for any $\alpha \in [0, 1]$ and $x \in I$, $k = 0, 1, 2, \dots$

Theorem 1.7 Let $g : I \rightarrow \mathbb{R}$ be a strictly convex function unbounded on I with a continuous derivative $g' : J \rightarrow \mathbb{R}$ such that $g \geq c$ for some constant $c > 0$. In case I is unbounded, we additionally require

$$\lim_{|t| \rightarrow \infty} \frac{g(t)}{|t|} = +\infty.$$

We assume that $C(I, g) \subset D$ and the following conditions **(G)** and **(H_g)** hold,

(G) $L_p(g_{n,x}^p)(x) \geq g_{n,x}^p(x) = g(x)$ for any $x \in J$, $n \geq 1$ and $p = 0, \dots, n$,

(H_g) $M_g(x) := \sup_{p \in \mathbb{N}} p |L_p(g)(x) - g(x)|$ is finite for every $x \in J$.

1. If $(L_n)_{n \geq 1}$ satisfies **(H₁)** with growth M_1 , then for all $f \in C(I, g)$, we have

$$(1.11) \quad \lim_{n \rightarrow \infty} L_{n,\lambda_n}(f) = f$$

pointwise, and the convergence is uniform on the sets where M_1 , M_g and g' are bounded.

2. If $(L_n)_{n \geq 1}$ satisfies $(\mathbf{H}_2)$ with growth M_2 , then there exists a constant $M > 0$, depending only on f , I and J , such that for every $n \geq 1$, $x \in J$, we have

$$(1.12) \quad \begin{aligned} |L_{n,\lambda_n}(f)(x) - f(x)| &\leq 2\omega\left(f, \sqrt{\frac{\lambda_n(x)M_2(x)}{n}}\right) + \\ &+ M \left[\frac{\lambda_n(x)M_g(x)}{n} + 2|g'(x)| \sqrt{\frac{M_2(x)\lambda_n(x)}{n}} \right]. \end{aligned}$$

Moreover, if also $(\mathbf{H}_1)$ holds with growth M_1 , then for $n \geq 1$, $x \in J$, we have

$$(1.13) \quad \begin{aligned} |L_{n,\lambda_n}(f)(x) - f(x)| &\leq 2\omega\left(f, \sqrt{\frac{\lambda_n(x)M_2(x)}{n}}\right) + \\ &+ M \frac{\lambda_n(x)M_g(x) + |g'(x)| M_1(x)(1 - (1 - \lambda_n(x))^n)}{n}. \end{aligned}$$

Proof. We start with some consequence of conditions $(\mathbf{G})$ and $(\mathbf{H}_g)$.

The identity

$$(1.14) \quad L_{n,\lambda_n}(f)(x) - f(x) = \sum_{p=0}^n \binom{n}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-p} (L_p(f_{p/n,x})(x) - f(x)),$$

and $(\mathbf{G})$ assures that $L_{n,\lambda_n}(g)(x) - g(x) \geq 0$.

The convexity of g yields

$$g_{\frac{p}{n},x}^p(t) - g(x) \leq \frac{p}{n}(g(t) - g(x)),$$

which with identity (1.14) and $(\mathbf{H}_g)$ implies

$$\begin{aligned} 0 \leq L_{n,\lambda_n}(g)(x) - g(x) &\leq \sum_{p=1}^n \binom{n}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-p} \frac{p}{n} (L_p(g)(x) - g(x)) \\ &\leq \frac{M_g(x)}{n} \sum_{p=1}^n \binom{n}{p} \lambda_n(x)^p (1 - \lambda_n(x))^{n-p} = \frac{M_g(x)\lambda_n(x)}{n}. \end{aligned}$$

1. Relation (1.10) together with the previous one, assure the convergence of L_{n,λ_n} on the “test” functions $\mathbf{1}$, e_1 and g . Now, using Theorem 4.1 in [6], we obtain the convergence (1.11).

2. In this case the relation (1.9) and (1.8) hold. Thus using Theorem 4.5 in [6] we get (1.12). Moreover, if even $(\mathbf{H}_1)$ holds, then (1.10) and again Theorem 4.5 in [6], yield (1.13). $\square$

Observe that condition $(\mathbf{G})$ is satisfied whenever L_n has the following property:

$$f \text{ is convex} \Rightarrow L_n(f) \geq f.$$

Another concrete condition implying $(\mathbf{G})$ is provided by the following lemma.

Lemma 1.8 *We assume that L_n preserves the affine function, that is $L_n(e_i) = e_i$ for $i = 0, 1$. For any convex function $f \in C^2(I) \cap D$, we have $L_n(f) \geq f$.*

Proof. Indeed, expanding f in Taylor series, we have

$$f(u) = f(x) + f'(x)(u - x) + \frac{f''(\eta)}{2}(u - x)^2 \geq f(x) + f'(x)(u - x),$$

and applying L_n to both side of the inequality, we have $L_n(f) \geq f$. $\square$

Actually, also condition $(\mathbf{H}_g)$ is easily checkable. For instance, it is satisfied if a Voronovskaja-type relation holds for L_n (see next example).

Example. Set $EXP(I) := \bigcup_{w>0} C(I, \exp(we_1) + \exp(-we_1))$, the space of continuous functions with exponentially growth. For any $n \geq 1$ let $L_n: EXP(I) \rightarrow C(J)$ be a linear positive operator defined by

$$L_n(f)(x) = \int_I W_n(x, t) f(t) dt,$$

where the kernel $W_n(x, t)$ is a generalized positive function. We assume that there exists a strictly positive function $p \in C(I)$, analytic on the inner of I , such that the relations

$$(1.15) \quad \begin{aligned} L_n(\mathbf{1})(x) &= 1, \\ \frac{\partial}{\partial x} L_n(f)(x) &= \frac{n}{p(x)} L_n(\psi_x f)(x), \end{aligned}$$

hold for any $f \in EXP(I)$ and $x \in J$. These operators are also referred as the *exponential operators* (for more details see [8, 9]). Operators satisfying these property are, for example, the Bernstein polynomials, the Szász-Mirakjan, Baskakov, Post-Widder and Weierstrass operators.

These operators realize approximation processes for functions in $EXP(I)$, and the following results hold (see [8, 9]).

Proposition 1.9 *For any $n \geq 1$, $x \in J$ and $f \in EXP(I) \cap C^2(I)$, the following assertions hold:*

1. $L_n(\mathbf{1})(x) = 1$,
2. $L_n(e_1)(x) = x$,
3. $L_n(e_2)(x) = x^2 + \frac{p(x)}{n}$,
4. $\lim_n n(L_n(f)(x) - f(x)) = \frac{1}{2}p(x)f''(x)$ uniformly on any compact subset of the interior of J .

Now, we can state the following theorem.

Theorem 1.10 *For any $f \in C(I)$, such that $|f(x)| \leq K(\exp(wx) + \exp(-wx))$, there exists a constant $M(f) > 0$, depending only on f , for which the estimate*

$$|L_{n,\lambda_n}(f)(x) - f(x)| \leq 2\omega\left(f, \sqrt{\frac{\lambda_n(x)p(x)}{n}}\right) + M(f)w^2 \frac{\lambda_n(x)p(x) \cosh(wx)}{n},$$

holds uniformly on compact subset of J .

Proof. By Proposition 1.9, $(L_n)_{n \geq 1}$ satisfies $(\mathbf{H}_2)$ with $M_2 = p$ and $(\mathbf{H}_1)$ with $M_1 = 0$. We set $g(x) = \exp(wx) + \exp(-wx)$. From Lemma 1.8 and 4. of Proposition 1.9, we deduce respectively that $(\mathbf{G})$ and $(\mathbf{H}_g)$ are satisfied. Thus, applying Theorem 1.7, we obtain the claim. $\square$

2 Voronovskaja-type results

Proposition 2.1 *Assume J compact, there exist $\alpha, \beta \in C(J)$, an even number $q > 2$ such that $e_k \in D$ for $k = 2, \dots, q$, and for every $x \in J$, there hold*

- (i) $\lim_{n \rightarrow \infty} nL_n(\psi_x)(x) = \beta(x)$,
- (ii) $\lim_{n \rightarrow \infty} nL_n(\psi_x^2)(x) = \alpha(x)$,
- (iii) $\lim_{n \rightarrow \infty} nL_n(\psi_x^q)(x) = 0$.

Moreover assume that the sequence $(\lambda_n)_{n \geq 1}$ converges uniformly to a function λ , and $\lambda_n(x) > 0$ for every $n \geq 1$ and $x \in J$. Then for every $f \in C^2(I) \cap D$, such that f'' is bounded, the convergence

$$(2.1) \quad \lim_{n \rightarrow \infty} n(L_{n,\lambda_n}(f)(x) - f(x)) = \lambda(x) \frac{\alpha(x)}{2} f''(x) + \beta(x) f'(x)$$

holds pointwise on J .

In particular, if $\lambda = 0$, then

$$\lim_{n \rightarrow \infty} n(L_{n,\lambda_n}(f)(x) - f(x)) = \beta(x) f'(x).$$

Proof. Let $f \in C^2(I) \cap D$ bounded on I and $x \in J$. Set $M := \max_I f''$. Then

$$f = f(x)\mathbf{1} + f'(x)\psi_x + \frac{f''(x)}{2}\psi_x^2 + \omega(x, \cdot)\psi_x^2,$$

where $\omega(x, y) = \frac{f''(\xi) - f''(x)}{2}$ with ξ between x and y . Hence

$$n(L_{n,\lambda_n}(f)(x) - f(x)) = f'(x)nL_{n,\lambda_n}(\psi_x)(x) + \frac{f''(x)}{2}nL_{n,\lambda_n}(\psi_x^2) + nL_{n,\lambda_n}(\omega(x, \cdot)\psi_x^2)(x).$$

By 1.2 in Lemma 1.1 and by Toeplitz theorem (see e.g. [11], Theorem (7.85)), as n tends to infinity, $nL_{n,\lambda_n}(\psi_x)(x)$ tends to $\beta(x)$ and $nL_{n,\lambda_n}(\psi_x^2)(x)$ tends to $\alpha(x)$. Moreover f'' is continuous in x , thus, for a fixed $\varepsilon > 0$ there exists $\delta > 0$ such that $|x - y| < \delta$ implies $|f''(x) - f''(y)| < \varepsilon$, for every $y \in I$.

Therefore, if $|x - y| < \delta$, then $|\omega(x, y)\psi_x^2(y)| \leq \varepsilon\psi_x^2(y)$, while, if $|x - y| \geq \delta$, then $|\omega(x, y)\psi_x^2(y)| \leq M\delta^{2-q}\psi_x^q(y)$. Hence

$$|nL_{n,\lambda_n}(\omega(x, \cdot)\psi_x^2)(x)| \leq \varepsilon nL_{n,\lambda_n}(\psi_x^2)(x) + M\delta^{2-q}nL_{n,\lambda_n}(\psi_x^q)(x).$$

Thus, by applying again Toeplitz theorem, we get $\lim_{n \rightarrow \infty} nL_{n,\lambda_n}(\omega(x, \cdot)\psi_x^2)(x) = 0$. $\square$

Example 1. The convergence in the Voronovskaja-formula (2.1) is uniform on J when $\lambda_n(x) \geq \lambda_0$ for every $n \geq 1$, $x \in J$, and the convergence in (i), (ii), (iii) are uniform on J , since in this case Toeplitz theorem yields the uniform convergence (see examples after Theorem (7.85) in [11]). But the convergence can be uniform even when $\lambda = 0$.

For this we discuss the case of Bernstein-Schurer operators $B_{n,1} : C([0, 2]) \rightarrow C([0, 1])$, defined for every $f \in C([0, 2])$ and $x \in [0, 1]$ by

$$B_{n,1}(f) := \sum_{k=0}^{n+1} \binom{n+1}{k} x^k (1-x)^{n+1-k} f\left(\frac{k}{n}\right)$$

(see e.g. 5.3.1 in [1]).

It is known that $B_{n,1}(\psi_x)(x) = \frac{1}{n}x$, and $B_{n,1}(\psi_x^2)(x) = \frac{1}{n}x(1-x) + \frac{x}{n^2}$. Simple calculations show that for every $x \in [0, 1]$, $B_{n,1}(\psi_x^4)(x) \leq \frac{M}{n^3}$, where M is a constant that does not depend on n nor on x .

Choose $\lambda_n = \frac{1}{n}$ for every $n \geq 1$. Then

$$B_{n,1,\lambda_n}(\psi_x)(x) = \frac{x}{n}, \quad B_{n,1,\lambda_n}(\psi_x^2)(x) = \frac{x}{n^2} + \frac{1}{n^2}x(1-x), \quad B_{n,1,\lambda_n}(\psi_x^4)(x) \leq \frac{M}{n^3}.$$

By reviewing the proof of 2.1, one concludes that for every $f \in C^2([0, 2])$,

$$\lim_{n \rightarrow \infty} n(B_{n,1,\lambda_n}(f) - f)(x) = xf'(x)$$

uniformly on $[0, 1]$.

Example 2. Let $(L_n)_{n \geq 1}$ be the sequence of exponential operators that have been considered in the example in the previous section. By combining (1.15) and 3. of Proposition 1.9, one has $\lim_n nL_n(\psi_x^4)(x) = 0$. Now, in the setting of Proposition 2.1, that is, assuming J compact, $\lambda_n > 0$ for any $n \geq 1$ and $\lambda_n \rightarrow \lambda$ uniformly, we have that, for every $f \in C^2(I) \cap EXP(I)$ such that $f'' \in C_b(I)$, there holds

$$\lim_n n(L_{n,\lambda_n}(f)(x) - f(x)) = \frac{\lambda(x)p(x)}{2}f''(x) \quad (x \in J).$$

3 Shape preserving properties

We investigate the behaviour of the sequence $(L_{n,\lambda_n})_{n \geq 1}$ when acting on convex, monotone or Hölder continuous functions. The results do not depend on the topological structure of the space D , but only on the positivity of the operators L_n and their shape preserving properties.

Proposition 3.1 *For every $n \geq 1$, let $\lambda_n = \lambda : J \rightarrow [0, 1]$. Assume that for every convex function $f \in D$, the sequence $(L_n(f))_{n \geq 1}$ is decreasing. Then $(L_{n,\lambda}(f))_{n \geq 1}$ is also decreasing for every convex function $f \in D$.*

Proof. Let $f \in D$ be a convex function and $x \in J$. We have to prove that the quantity $Q := L_{n+1,\lambda}(f)(x) - L_{n,\lambda}(f)(x) = L_{n+1,\lambda}(f)(x) - L_{n,\lambda}(f)(x)(1 - \lambda(x) + \lambda(x))$ is not positive. For sake of shortness, in the sequel of this proof we avoid to indicate the dependence from x of λ , f and L_p . By computation one gets

$$\begin{aligned} Q &= \sum_{p=0}^{n+1} \binom{n+1}{p} \lambda^p (1-\lambda)^{n+1-p} L_p(f_{p/(n+1),x}) - \sum_{p=0}^n \binom{n}{p} \lambda^{p+1} (1-\lambda)^{n-p} L_p(f_{p/n,x}) + \\ &\quad - \sum_{p=0}^n \binom{n}{p} \lambda^p (1-\lambda)^{n+1-p} L_p(f_{p/n,x}) = \\ &= (1-\lambda)^{n+1} f + \lambda^{n+1} L_{n+1}(f) - (1-\lambda)^{n+1} f - \lambda^{n+1} L_n(f) + \\ &\quad + \left[(n+1) L_1(f_{1/(n+1),x}) - f - n L_1(f_{1/n,x}) \right] \lambda (1-\lambda)^n + \sum_{p=2}^n \lambda^p (1-\lambda)^{n+1-p} \times \\ &\quad \times \left[\binom{n+1}{p} L_p(f_{p/(n+1),x}) - \binom{n}{p-1} L_{p-1}(f_{(p-1)/n,x}) - \binom{n}{p} L_p(f_{p/n,x}) \right]. \end{aligned}$$

For every $n \geq 1$ and $p = 0, \dots, n-1$ the function $f_{p/n,x}$ is convex, therefore $L_p(f_{p/n,x}) \geq L_{p+1}(f_{p/n,x})$. By the positivity of the operators L_p and the convexity of f , we obtain

$$\begin{aligned} Q &\leq \lambda (1-\lambda)^n \left[(n+1) \left(\frac{n}{n+1} L_1(f_{1/n,x}) + \frac{1}{n+1} f \right) - f - n L_1(f_{1/n,x}) \right] + \\ &\quad + \sum_{p=2}^n \left[\binom{n+1}{p} L_p(f_{p/(n+1),x}) - \binom{n}{p-1} L_{p-1}(f_{(p-1)/n,x}) - \binom{n}{p} L_p(f_{p/n,x}) \right] \lambda^p (1-\lambda)^{n+1-p} \\ &= \sum_{p=2}^n L_p \left[\binom{n+1}{p} f_{p/(n+1),x} - \binom{n}{p-1} f_{(p-1)/n,x} - \binom{n}{p} f_{p/n,x} \right] \lambda^p (1-\lambda)^{n+1-p}. \end{aligned}$$

From the convexity of f and positivity of L_p , we get $Q \leq 0$. □

Corollary 3.2 *If for every convex function $f \in D$ and $n \geq 1$,*

$$f \leq L_{n+1}(f) \leq L_n(f),$$

then for every convex function $f \in D$ and $n \geq 1$, we have

$$f \leq L_{n+1,\lambda}(f) \leq L_{n,\lambda}(f) \leq \lambda L_1(f) + (1-\lambda)(f) \leq L_1(f).$$

Proof. Let $f \in D$ be convex, then $f_{p/n,x}$ is convex for every $n \geq 1$ and $p = 0, \dots, n$. Hence for every $t \in J$

$$L_p(f_{p/n,x})(t) \geq f_{p/n,x}(t).$$

In particular, $L_p(f_{p/n,x})(x) \geq f(x)$, hence $L_{n,\lambda}(f)(x) \geq \sum_{p=0}^n \binom{n}{p} \lambda(x)^p (1 - \lambda(x))^{n-p} f(x) = f(x)$. The other inequalities follow by observing that $L_{1,\lambda} = \lambda L_1 + (1 - \lambda)Id_D$ and that $f \leq L_1(f)$. $\square$

Consider the following subsets of $F(I)$:

$$\mathbf{M}_+(I) := \{f : I \rightarrow \mathbb{R} : f \text{ is increasing}\},$$

$$\mathbf{Cx}(I) := \{f : I \rightarrow \mathbb{R} : f \text{ is convex}\},$$

$$\mathbf{Lip}_K(\alpha)(I) := \{f : I \rightarrow \mathbb{R} : |f(x) - f(y)| \leq K|x - y|^\alpha \ (x, y \in I)\} \ (K > 0, \alpha > 0).$$

Proposition 3.3 *Let $\lambda_n \in [0, 1]$ be a constant.*

1. *If $L_p(\mathbf{M}_+(I) \cap D) \subseteq \mathbf{M}_+(J)$ for every $p = 1, \dots, n$, then $L_{n,\lambda_n}(\mathbf{M}_+(I) \cap D) \subseteq \mathbf{M}_+(J)$.*
2. *If $L_p(\mathbf{M}_+(I) \cap D) \subseteq \mathbf{M}_+(J)$ and $L_p(\mathbf{Cx}(I) \cap D) \subseteq \mathbf{Cx}(J)$ for every $p = 1, \dots, n$, then $L_{n,\lambda_n}(\mathbf{Cx}(I) \cap D) \subseteq \mathbf{Cx}(J)$.*
3. *If $L_p(\mathbf{Lip}_K(\alpha)(I) \cap D) \subseteq \mathbf{Lip}_H(\alpha)(J)$ for every $p = 1, \dots, n$, then $L_{n,\lambda_n}(\mathbf{Lip}_K(\alpha)(I) \cap D) \subseteq \mathbf{Lip}_{K+H}(\alpha)(J)$. In particular Lipschitz continuous functions are preserved.*

Proof. 1. See the proof of Proposition 2.2 in [5].

2. Let $f \in \mathbf{Cx}(I) \cap D$ and $x, y \in I$, $x < y$. As already stated in Remark 1.6 $f_{p/n,x} \in \mathbf{Cx}(I)$ for every $p = 0, \dots, n$ and $x \in I$. By following the idea of the proof of Proposition 2.3 in [5], one gets

$$L_p(f_{p/n,y})(y) + L_p(f_{p/n,y})(x) \leq L_p(f_{p/n,x})(x) + L_p(f_{p/n,y})(y).$$

Now, since the operator L_p is positive and preserves the convexity, we get for every $t \in [0, 1]$,

$$\begin{aligned} L_p(f_{p/n,tx+(1-t)y})(tx + (1-t)y) &\leq \\ &\leq tL_p(f_{p/n,tx+(1-t)y})(x) + (1-t)L_p(f_{p/n,tx+(1-t)y})(y) \\ &\leq t^2L_p(f_{p/n,x})(x) + t(1-t)L_p(f_{p/n,y})(y) + \\ &\quad + (1-t)tL_p(f_{p/n,x})(x) + (1-t)^2L_p(f_{p/n,y})(y) \\ &= tL_p(f_{p/n,x})(x) + (1-t)L_p(f_{p/n,y})(y). \end{aligned}$$

Thus, the function $z \rightarrow L_p(f_{p/n,z})(z)$ is convex. It follows immediately that $L_{n,\lambda_n}(f)$ is convex too.

3. Let $f \in \mathbf{Lip}_K(\alpha)(I) \cap D$ and $x, y \in I$. Then $f_{p/n,x} \in \mathbf{Lip}_{K(p/n)^\alpha}(\alpha)(I)$ and hence

$$\begin{aligned} |L_{n,\lambda_n}(f)(x) - L_{n,\lambda_n}(f)(y)| &\leq \sum_{p=0}^n \binom{n}{p} \lambda_n^p (1 - \lambda_n)^{n-p} \times \\ &\quad \times [|L_p(f_{p/n,x})(x) - L_p(f_{p/n,x})(y)| + |L_p(f_{p/n,y})(y) - L_p(f_{p/n,x})(y)|] \\ &\leq \sum_{p=0}^n \binom{n}{p} \lambda_n^p (1 - \lambda_n)^{n-p} \left[K \left(\frac{p}{n} \right)^\alpha + HL_p \left(\left(1 - \frac{p}{n} \right)^\alpha |x - y|^\alpha \cdot \mathbf{1} \right)(y) \right] \\ &\leq (H + K)|x - y|^\alpha, \end{aligned}$$

that is the claim. $\square$

Remark 3.4 *If λ_n is not a constant, the previous results do not hold in general. In [5], there are several counterexamples in the case of Szász-Mirakjan operators.*

Finally, we do some remarks about the monotonicity of the operators L_{n,λ_n} with respect to the function λ_n . If $\lambda \leq \alpha$, in general we do not have $L_{n,\lambda}(f) \leq L_{n,\alpha}(f)$, even when α and λ are constants and f is convex. For a counterexample consider $\lambda = \frac{1}{2}$, $\alpha = 1$, L_n be the n -th Kantorovich operator on $[0, 1]$ and $f = e_1$ (see [1], 5.3.7).

Proposition 3.5 *Let $\alpha_n, \lambda_n : J \rightarrow [0, 1]$ be functions such that $\alpha_n \leq \lambda_n$. If $f \in D$ and $x \in J$ satisfy $L_p(f_{p/n,x})(x) \leq L_{p+1}(f_{(p+1)/n,x})(x)$ for every $p = 0, \dots, n-1$, then $L_{n,\alpha_n}(f)(x) \leq L_{n,\lambda_n}(f)(x)$.*

Proof. Let $f \in D$ and $x \in J$ satisfying the assumptions. The function

$$\phi : s \in [0, 1] \rightarrow \sum_{p=0}^n \binom{n}{p} s^p (1-s)^{n-p} L_p(f_{p/n,x})(x),$$

has derivative

$$\phi'(s) = n \sum_{p=0}^n \binom{n-1}{p} s^p (1-s)^{n-p-1} (L_{p+1}(f_{(p+1)/n,x})(x) - L_p(f_{p/n,x})(x)) \quad (s \in [0, 1]),$$

hence it is increasing. In particular $L_{n,\alpha_n}(f)(x) = \phi(\alpha_n(x)) \leq \phi(\lambda_n(x)) = L_{n,\lambda_n}(f)(x)$. $\square$

Remark 3.6 *If for every $x \in J$, there exists a regular finite Borel measure γ_x on I such that*

$$L_p(f)(x) = \int_{I^p} f \left(\frac{x_1 + \dots + x_p}{p} \right) d\gamma_x(x_1) \cdots d\gamma_x(x_p) \quad (f \in D),$$

and $f|_J \leq L_1(f)$ for every convex function f , then $L_p(f_{p/n,x})(x) \leq L_{p+1}(f_{(p+1)/n,x})(x)$ for every convex function f , $x \in J$, $n \geq 1$ and $p = 1, \dots, n-1$.

Indeed, let $x_0, x \in J$, then for every $\alpha \in [0, 1]$, we have

$$f(\alpha x_0 + (1-\alpha)x) = f_{\alpha,x_0}(x) \leq L_1(f_{\alpha,x})(x) = \int_I f(\alpha x_0 + (1-\alpha)t) d\gamma_x(t).$$

Since $\frac{x_1 + \dots + x_p + x}{p+1} \in I$, whenever $x_1, \dots, x_p, x \in I$, we get

$$\begin{aligned} L_p(f_{p/n,x})(x) &= \int_{I^p} f \left(\frac{x_1 + \dots + x_p + x}{p+1} \frac{p+1}{n} + \frac{(n-p-1)x}{n} \right) d\gamma_x(x_1) \cdots d\gamma_x(x_p) \\ &\leq \int_{I^p} \left(\int_I f \left(\frac{x_1 + \dots + x_p + x_{p+1}}{n} + \frac{(n-p-1)x}{n} \right) d\gamma_x(x_{p+1}) \right) d\gamma_x(x_1) \cdots d\gamma_x(x_p) \\ &= L_{p+1}(f_{(p+1)/n,x})(x). \end{aligned}$$

Particular cases of this situation are Bernstein-Schnabl operators (see [1]), Szász-Mirakjan and Baskakov operators on $[0, +\infty[$ and their modification of Lototskij-Schnabl type (see [1, 2, 3]).

References

- [1] F. ALTOMARE, M. CAMPITI, *Korovkin-type Approximation Theory and its Applications*, De Gruyter Stud. in Math. **17**, Walter de Gruyter, Berlin-New York, 1994
- [2] F. ALTOMARE, I. CARBONE, On some degenerate differential operators on weighted function spaces, *J. Math. Analysis Appl.* **213** (1997), 308–333
- [3] F. ALTOMARE, E.M. MANGINO, On a generalization of Baskakov operators, *Rev. Roum. Matem. Pures Appl.* XLIV, 5-6 (1999), 683–705
- [4] F. ALTOMARE, E.M. MANGINO, On a class of elliptic-parabolic equations on unbounded intervals, to appear in *Positivity*, 2000
- [5] I. CARBONE, Shape preserving properties of some positive linear operators on unbounded intervals, *J. Approx. Theory* **92** (1998), 140–156
- [6] L. D’AMBROSIO, Approximation of $\ast$ -weak-to-norm mappings, Preprint SISSA 75/2000/M (2000). Submitted.
- [7] Z. DITZIAN, Convergence of Sequences of Linear Positive Operators: Remarks and Applications, *J. of Approx. Theory* **14** (1975), 296–301.
- [8] M.E.H. ISMAIL, C.P. MAY, On a Family of Approximation Operators, *J. Math. Anal. Appl.* **63** (1978), 446–462.
- [9] C.P. MAY, Saturation and Inverse Theorems for Combinations of a Class of Exponential-Type Operators, *Can. J. Math.* **28** (1976), 1224–1250.
- [10] S.-Y. SHAW, Approximation of unbounded functions and applications to representations of semigroups, *J. Approx. Theory* **28** (1980), 238–259.
- [11] K.R. STROMBERG, *Introduction to Classical Real Analysis*, Wadsworth International Group, Belmont, California (1981)
- [12] V. TOTIK, Uniform approximation by positive operators on infinite intervals, *Anal. Math.* **10** (1984), 163–182

Author’s address:

LORENZO D’AMBROSIO
 SISSA - ISAS v. Beirut, 2-4
 I-34014 TRIESTE ITALY
 E-MAIL:dambros@sissa.it

ELISABETTA MANGINO
 DIPARTIMENTO DI MATEMATICA “E. DE GIORGI”
 UNIVERSITA’ DEGLI STUDI DI LECCE
 I-73100 LECCE
 E-MAIL:elisabetta.mangino@unile.it